

Chest Wall Injury

Chest wall injury guidelines

- [Care of Patients with Rib Fractures](#)
- [Care of Patients with Rib Fractures](#)
- [Surgical Stabilization of Rib Fractures \(SSRF or Rib Plating\)](#)

Care of Patients with Rib Fractures

Purpose:

Rib fractures occur in approximately 10% of patients with traumatic injury. They are associated with a greater injury burden especially when coupled with head, extremity, abdominal, and blunt cardiac injury. Mortality rates increase with the number of fractured ribs; 5.8% for a single rib fracture to 34.4% mortality with 8 or more rib fractures. Flail chest and pulmonary contusion also increase mortality. Rib fractures are associated with multiple pulmonary complications including pneumonia, adult respiratory distress syndrome, and pneumothorax. Rib fractures are also associated with an increased ICU length of stay, hospital length of stays, and ventilator days. The purpose of this guideline is to standardize our approach to the management of traumatic rib fractures.

Admission Criteria:

Admit to unit based on age, injury burden, degree of pulmonary compromise, comorbidities, and at trauma attending's discretion.

1. Consider admission to ICU if:
 - Mechanical Ventilation
 - Age >60
 - 4 or more rib fractures
 - Lung parenchymal abnormality or contusion
 - Flail segment
 - Volume expansion protocol needed more frequently than every 2 hours
 - Incentive spirometry <1000 cc
 - COPD, Home Oxygen Use, Current Tobacco User, Antiplatelet Use².
2. Consider admission to Step-Down Critical Care Unit if:
 - <3 rib fractures
 - Age >45 with rib fractures and flail segment or sternal fracture
 - Oxygen requirement \geq 5L/min nasal cannula
 - Volume expansion protocol needed every 2-3 hours
 - Incentive spirometry 1000-1500 cc
3. Consider Floor admission if:
 - Pain control is adequate
 - Incentive spirometry >1500 cc

Initial Management:

1. Consult to Respiratory Therapy for Lung Volume Expansion (if no pneumothorax).
2. Continuous pulse oximetry.
3. Incentive spirometry for 10 times per hour while awake.
4. Supplemental oxygen as needed to maintain SpO₂ >90% (or >88% in patients with known history of COPD).
5. CXR PA portable every morning for 3 days (+/- days based on clinical judgment).
6. Physical therapy consult for early mobilization. When cleared, patient should ambulate three times a day at minimum.
7. Judicious use of intravenous fluids. Avoid boluses. If boluses are indicated utilize small boluses. If unresponsive to 2 boluses, notify trauma attending.
8. Multimodal Pain Management:
 - PCA or hourly prn IV pain medication.
 - Consult to APS for epidural or paravertebral block if not contraindicated
 - **Contraindications for an epidural include:** platelets <80K, infection at site of insertion, epidural or spinal cord hematoma, INR >1.2, prophylactic LMWH within 12 hours or therapeutic dose within 24 hours, hemodynamic instability.
 - **Contraindications for a paravertebral block include:** platelets <80K, infection at site of insertion, INR >1.5, transverse process fracture in proximity to level of insertion.
 - Lidocaine patch over rib fractures
 - Tylenol 1000 mg PO every 6 hours scheduled + Flexeril 10 mg PO every 8 hours scheduled + Oxycodone Immediate Release 5-15 mg PO every 4 hours as needed PRN
 - Add ibuprofen 800 mg PO every 8 hours scheduled if not contraindicated due to age, renal function or bleeding risk; strongly consider a COX-2 inhibitor is ibuprofen contraindicated.

Non-invasive mechanical ventilation (BiPAP or CPAP):

Should only be used if the patient is normally on this treatment prior to injury.

1. BiPAP/CPAP is rarely appropriate for patients with chest injuries and progressive respiratory distress. Early intubation in these patients is more appropriate.
2. BiPAP should be used for reversible ventilation issues such as hypercarbia, COPD exacerbation, and/or pulmonary edema.
3. BiPAP is a bridge to allow time for interventions (e.g. Lasix administration) to be performed that may prevent intubation.
4. BiPAP should only be used as a short term option, ideally no more than 6 hours.
5. Monitor the patient closely while on BiPAP for further respiratory decline.
6. If respiratory status does not improve within 6 hours or less, consider intubation.

Surgical Stabilization of Rib Fractures (Rib Plating):

Consider rib plating in the following clinical situations: (see Trauma Policy PRO06 Surgical Stabilization of Rib Fractures):

1. Non-intubated patients with respiratory insufficiency due to pain despite continuous epidural/paravertebral anesthesia.
2. Intubated patients with flail chest who fail to wean from ventilator.
3. Patients with extensive anterolateral flail chest and progressive displacement of fractured ribs.
4. Patients who require thoracotomy due to associated intra-thoracic injury.
5. Painful nonunion.
6. Patient complaints of painful movement of ribs.

References:

1. Carver, T., Milia, D, Somberg, C., Brasel, K., & Paul, J. (2015). Vital capacity helps predict pulmonary complications after rib fractures. *Journal of Trauma Acute Care Surgery*, 79(3), 413-416. doi: 10.1097/TA.0000000000000744
2. Chen, J., Jeremitsky, E., Philp, R. Fry, W., & Smith R. (2014). A chest trauma scoring system to predict outcomes. *Journal of Surgery*, 156(4), 988-994. doi: 10.1016/j.surg.2014.06.045
3. Gonzalez, K., Ghneim, M., Kang, F., Jupiter, D., Davis, M., Regner, J. (2015). A pilot single-institution predictive model to guide rib fracture management in elderly patients. *Journal of Trauma Acute Care Surgery*, 78(5), 970-975. doi: 10.1097/TA.0000000000000619
4. Leininger, S. (2017). Rib fracture protocol advancing the care of the elderly patient. *Critical Care Nursing*, 40(1).
5. Mastroianni, S. Implementing a rib fracture management pathway and PIC scoring tool to reduce ICU readmissions. San Francisco, CA: University of San Francisco Scholarship Repository; May 22, 2015, Spring.
6. Sahr, S., Webb, M., Hackett Renner, C, Sokol, R., & Swegle, J. (2013). Implementation of a rib fracture triage protocol in elderly trauma patients. *Journal of Trauma Nursing*, 20(4), 172-175. doi: 10.1097/JTN.0000000000000008
7. Simon, B., Ebert, J., Bokhari, F., Capella, J., Emohoff, T., Hayward, T., Rodriguez, A., & Smith, L. (2012). Management of pulmonary contusion and flail chest: An Eastern Association for the Surgery of Trauma practice management guidelines. *Journal of Trauma Acute Care Surgery*, 73(5), S351-S361. doi: 10.1097/TA.0b013e31827019fd
8. Witt, C., & Bulger, E. (2017). Comprehensive approach to the management of the patient with multiple rib fractures: a review and introduction of a bundled rib fracture management protocol. *Trauma Surgery & Acute Care Open*, 2(1), 1-7. doi: 10.1136/tsaco-2016-000064

DATE: 7/6/17

Care of Patients with Rib Fractures

Purpose

Rib fractures occur in approximately 10% of patients with traumatic injury. They are associated with greater injury burden especially when coupled with head, extremity, abdominal and blunt cardiac injury. Mortality rates increase with the number of fractured ribs (5.8% for a single rib to 34.4% mortality with 8 or more rib fractures). Flail chest and pulmonary contusion also increase mortality. Rib fractures are associated with multiple pulmonary complications including pneumonia, adult respiratory distress syndrome (ARDS) and pneumothorax. Rib fractures are also associated with an increased ICU length of stay (LOS), hospital LOS, and ventilator days. The purpose of this guideline is to standardize our approach to the management of traumatic rib fractures.

Admission Criteria:

Admit to unit based on age, injury burden, degree of pulmonary compromise, comorbidities, and trauma attending discretion.

1. Consider admission to ICU if:
 - mechanical ventilation
 - age > 60 yrs
 - 4 or more rib fractures
 - lung parenchymal abnormality or contusion
 - flail segment
 - volume expansion protocol needed more frequently than every 2 hours
 - incentive spirometry <1000 cc.
 - COPD, home O2 use, current tobacco user, current antiplatelet use
2. Consider admission to STEP DOWN Unit if:
 - <3 rib fractures
 - age >45 yrs with rib fractures and flail segment or sternal fracture
 - O2 requirement greater than or equal to 5L nasal canula
 - volume expansion protocol needed every 2-3 hrs
 - incentive spirometry 1000-1500 cc
3. Consider admission to FLOOR if:
 - pain control is adequate
 - incentive spirometry >1500 cc

Initial Management:

1. Consult to respiratory therapy for "Lung Volume Expansion" (if no pneumothorax)
2. Continuous pulse oximetry
3. Incentive spirometry for 10 times/hr while awake
4. Supplemental oxygen as needed to maintain SpO₂>90% (or >88% in patients with known history of COPD).
5. Chest X-ray (portable) every morning x 3 days (+/1 days based on clinical judgement)
6. Physical therapy consult for early mobilization. When cleared, patient should ambulate 3x daily at minimum.
7. Judicious use of intravenous fluids. Avoid boluses if possible and, if boluses are indicated, utilize small boluses. If unresponsive to 2 boluses, notify trauma attending.
8. Multimodality pain management:
 - PCA or hourly PRN IV pain medication
 - Consult APS for epidural or paravertebral block if not contraindicated.
 - contraindications for an epidural include: platelets <80K, infection at site of insertion, epidural or spinal cord hematoma, INR >1.2, prophylactic LMWH within 12 hrs or therapeutic dose within 24 hrs, hemodynamic instability.
 - contraindications to a paravertebral block include: platelets <80k, infection at site of insertion, INR <1.5, transverse process fractures in proximity to level of insertion.
 - Lidocaine patch over rib fractures
 - Tylenol 1000 mg PO every 6 hrs scheduled + Flexeril 10 mg PO every 8 hrs scheduled + Oxycodone immediate release 5-15mg PO every 4 hrs as needed (PRN).
 - Add ibuprofen 800 mg PO every 8 hrs scheduled if not contraindicated due to age, renal function, or bleeding risk; strongly consider a COX-2 inhibitor if ibuprofen is contraindicated.

Non-invasive mechanical ventilation (BiPAP or CPAP):

Should only be used if the patient is normally on this treatment prior to injury.

1. BiPAP/CPAP is rarely appropriate for patients with chest injuries and progressive respiratory distress. Early intubation in these patients is more appropriate.
2. BiPAP should be used for reversible ventilation issues such as hypercarbia, COPD exacerbation, and/or pulmonary edema.
3. BiPAP is a bridge to all time for interventions (e.g. Lasix administration for volume overload) to be performed that may prevent intubation.
4. BiPAP should only be used as a short term option, ideally no more than 6 hours.
5. Monitor the patient closely while on BiPAP for further respiratory decline.
6. If respiratory status does not improve within 6 hours or less, consider intubation.

Surgical Stabilization of Rib Fractures (Rib Plating):

Consider rib plating in the following clinical situations: (see Trauma Policy PRO06 Surgical Stabilization of Rib Fractures):

1. Non-intubated patients with respiratory insufficiency due to pain despite continuous epidural/paravertebral anesthesia and use of multi-modality pain regimen.
2. Intubated patients with flail chest who fail to wean from ventilator.
3. Patients with extensive anterolateral flail chest and progressive displacement of fractured ribs.
4. Patients who require thoracotomy due to associated intra-thoracic injury.
5. Painful nonunion
6. Patient complaints of painful movement of ribs (popping, clicking).

References

1. Carver T, Milia D, Somberg C, Brasel K, Paul J. Vital capacity helps predict pulmonary complications after rib fractures. *J Trauma Acute Care Surg.* 2015;79(3):413-416.
2. Chen J, Jeremitsky E, Philp R, Fry W, Smith R. A chest trauma scoring system to predict outcomes. *J Surg.* 2014;156(4): 988-994.
3. Gonzalez K, Ghneim M, Kang F, Jupiter D, Davis M, Regner J. A pilot single-institution predictive model to guide rib fracture management in elderly patients. *J Trauma Acute Care Surg.* 2015; 78(5):970-975.
4. Leininger S, Rib fracture protocol advancing the care of the elderly patient. *Crit Care Nursing.* 2017;40(1).
5. Mastroianni S. Implementing a rib fracture management pathway and PIC scoring tool to reduce ICU readmissions. San Francisco, CA: University of San Francisco Scholarship Repository; May 22, 2015, Spring.
6. Sahr S, Webb M, Hackett Renner C, Sokol R, Swegle J. Implementation of a rib fracture triage protocol in elderly trauma patients. *J Trauma Nursing.* 2013;20(4):172-175.
7. Simon B, Ebert J, Bkhari F, Capella J, Emohoff T, Hayward T, Rodriguez A, Smith L. Management of pulmonary contusion and flail chest. An Eastern Association for the Surgery of Trauma practice management guidelines. *J Trauma Acute Care Surg.* 2012;73(5):S351-S361.
8. Witt C, Bulger E. Comprehensive approach to the management of the patient with multiple rib fractures: A review and introduction of a bundled rib fracture management protocol. *Trauma Surg and Acute Care Open.* 2017;2(1): 1-7.

Author(s)

Trauma Leadership

Last Reviewed

July, 2017

Surgical Stabilization of Rib Fractures (SSRF or Rib Plating)

Purpose

Surgical Stabilization of rib fractures should be considered in patients with flail chest, flail sternum, and painful rib fractures associated with movement that have been refractory to conventional pain management in order to improve morbidity and mortality.

Indications

1. Non-ventilated patients:
 - Chest wall instability
 - 3 or more segmental rib fractures (flail chest)
 - 3 or more bi-cortically displaced/offset rib fractures
 - clinical findings of paradoxical motion
 - instability or "clicking" on palpation of chest wall or as reported by the patient
 - 3 or more displaced rib fractures
 - with displacement of >50% the rib width AND 2 or more pulmonary physiological derangements.
2. Ventilated patients:
 - Chest wall instability
 - 3 or more segmental rib fractures (flail chest)
 - 3 or more bi-cortically displaced/offset rib fractures
 - clinical findings of paradoxical motion
 - instability or "clicking" on palpation of chest wall or as reported by the patient
 - Failure to wean from ventilator

Contraindications

1. Absolute
 - shock/ongoing resuscitation
 - severe traumatic brain injury
 - acute myocardial infarction

- fractures outside of ribs 3-10

2. Relative

- Age <18 yrs
- Age >80 yrs
- unstable spine injury
- empyema
- history of chest wall radiation
- mild to moderate traumatic brain injury

Timing


1. Non-ventilated patients

- when feasible, less than 24 hrs is optimal
- should be performed within 72 hours of injury
- SSRF should be delayed in the face of higher priority injuries

2. Ventilated patients

- earliest feasible time for flail indication
- should be performed within 72 hrs of injury for non-flail indications.
- SSRF should be delayed in the face of higher priority injuries.

SSRF ALGORITHM



DEFINITIONS OF TERMS

SEVERE TBI

- ANY GCS <8
- SIGNS OF INTRACRANIAL HYPERTENSION

RELATIVE CONTRAINDICATIONS

- AGE <18 YEARS
- AGE >80 YEARS
- UNSTABLE SPINE INJURY
- EMPHYSEMA
- PRIOR CHEST WALL RADIATION
- MILD/MODERATE TBI

CHEST WALL INSTABILITY

FLAIL SEGMENT

- 3+ IPSILATERAL CONSECUTIVE RIBS WITH FRACTURES IN 2 LOCATIONS
- CLINICAL FINDING OF PARADOXICAL MOTION

OFFSET FRACTURES

- 3+ IPSILATERAL RIB FRACTURES WITH DISPLACEMENT OF 100% OF RIB WIDTH ON AXIAL CT

INSTABILITY OR "CLICKING" ON PALPATION OR REPORTED BY THE PATIENT

3+ > 50% DISPLACEMENT

- THREE IPSILATERAL CONSECUTIVE OR NON-CONSECUTIVE RIBS EACH WITH A FRACTURE DISPLACED 50% OF THE RIB WIDTH ON AXIAL CT

PULMONARY DERANGEMENTS

- RESPIRATORY RATE >20
- INCENTIVE SPIROMETRY <50% OF PREDICTED
- NUMERICAL PAIN SCORE >5/10
- POOR COUGH

FAILURE TO WEAN

MUST BE CLINICALLY DETERMINED TO BE RELATED TO THE RIB FRACTURES

UNABLE TO PROGRESS TO SPONTANEOUS BREATHING TRIAL AFTER 48 HOURS

ABLE TO OBTAIN SPONTANEOUS BREATHING TRIAL FOR 60 MINUTES BUT DEVELOPS ≥2 OF THE FOLLOWING

- INCREASED RESP. RATE >35
- INCREASED HEART RATE >140
- OXYGEN SATURATION <90%
- RSSI >105
- ANXIETY
- DIAPHORESIS
- AGITATION

OF NOTE: VENTILATOR WEANING SHOULD BE AT THE DISCRETION OF THE TREATING BEDSIDE PHYSICIAN.

HIGHER PRIORITY INJURIES

- PRE-OPERATIVE SPINAL INJURY
- OPEN ABDOMEN
- SIGNIFICANT VASCULAR TRAUMA
- PELVIC EXTERNAL FIXATION

References

1. Bauman, ZM., Grams, B., Yanala, U., Shostrom, V., Waibel, B., Evans, CH., Cemaj, S., & Schlitzkus, L. (2020). Rib fracture displacement worsens over time. *European Journal of Trauma and Emergency Surgery*. 2020 March 27. doi: 10.1007/s00068-020-01353-w
2. Cataneo, AJ., Cataneo, DC., de Oliveria, FH., Arruda, KA., El Dib, R., & de Oliveira Carvalho, PE. (2015). Surgical versus nonsurgical interventions for flail chest. *Cochrane Database of Systematic Reviews*, 2015(7), 1-30. doi: 10.1002/1465158.CD009919.pub2.
3. DeFreest, L., Tafen, M., Bhakta, A., Ata, A., Martone, S., Glotzer, O., Krautsak, K., Rosati, C., Stain, S., & Bonville, D. (2015). Open reduction and internal fixation of rib fractures in polytrauma patients with flail chest. *American Journal of Surgery*, 211(4), 761-767. doi: 10.1016/j.amjsurg.2015.11.014.
4. Kasotakis, G., Hasenboehler, EA., Streib, EW., Patel, N., Patel, MB., Alarcon, L., Bosarge, PL., Love, J., Haut, ER., & Como, JJ. (2017). Operative fixation of rib fractures after blunt trauma: A practice management guideline from the Eastern Association for the Surgery of Trauma. *The Journal of Trauma and Acute Care Surgery*, 82(3), 618-626. doi: 10.1097/TA.0000000000001350
5. Majercik, S., Vijayakumar, S., Olsen, G., Gardner, S., Granger, SR., Van Boerum, DH., & White, TW. (2015). Surgical stabilization of severe rib fractures decreased incidence of retained hemothorax and empyema. *American Journal of Surgery*, 210(8), 1112-1116. doi: 10.1016/j.amjsurg.2015.08.008.
6. Pieracci, FM., Coleman, J., Ali-Osman, F., Mangram, A., Majercik, S., White, TW., Jeremitsky, E., & Doben, AR. (2018). A multicenter evaluation of the optimal timing of surgical stabilization of rib fractures. *Journal of Trauma and Acute Care Surgery*, 84(1), 1-10. doi: 10.1097/TA.0000000000001729
7. Pieracci, FM., Lin, Y., Rodil, M., Synder, M., Herbert, B., Tran, DK., Stoval, RT., Johnson, JL., Biffi, WL., Barnett, CC., Cothren-Burlew, C., Fox, C., Jurkovich, GJ., & Moore, EE. (2018). A prospective, controlled clinical evaluation of surgical stabilization of severe rib fractures. *Journal of Trauma and Acute Care Surgery*, 80(2), 187-194. doi: 10.1097/TA.0000000000000925.
8. Pieracci, FM., Majercik, S., Ali-Osman, F., Ang, D., Doben, A., Edwards, JG., French, B., Gasparri, M., Marasco, S., Minshall, C., Sarani, B., Tisol, W., VanBoerum, DH., & White, TW. (2017). Consensus statement: Surgical stabilization of rib fractures rib fracture colloquium clinical practice guidelines. *Internal Journal of the Care of the Injured*, 48(2), 307-321. doi: 10.1016/j.injury.2016.11.026.
9. Witt, C., & Bulger, E. (2017). Comprehensive approach to the management of the patient with multiple rib fractures: a review and introduction of a bundled rib fracture management protocol. *Trauma Surgery & Acute Care Open*, 2(1), 1-7. doi: 10.1136/tsaco-2016-000064.
10. Wada, T., Yasuaga, H., Inokuchi, R., Matsui, H., Matsubara, T., Ueda, Y., Gunshin, M., Ishii, T., Doi, K., Kitsuta, Y., Nakajima, S., Fushimi, K., & Yahagi, N. (2015). Effectiveness of surgical rib fixation on prolonged mechanical ventilation in patients with traumatic rib fractures: A propensity score-matched analysis. *Journal of Critical Care*, 30(6), 1227-231. doi: 10.1016/j.jccr.2015.07.027.

Author(s)

Zachary Bauman, DO

Last Updated

May, 2020