

Surgical Stabilization of Rib Fractures (SSRF or Rib Plating)

Purpose

Surgical Stabilization of rib fractures should be considered in patients with flail chest, flail sternum, and painful rib fractures associated with movement that have been refractory to conventional pain management in order to improve morbidity and mortality.

Indications

1. Non-ventilated patients:
 - Chest wall instability
 - 3 or more segmental rib fractures (flail chest)
 - 3 or more bi-cortically displaced/offset rib fractures
 - clinical findings of paradoxical motion
 - instability or "clicking" on palpation of chest wall or as reported by the patient
 - 3 or more displaced rib fractures
 - with displacement of >50% the rib width AND 2 or more pulmonary physiological derangements.
2. Ventilated patients:
 - Chest wall instability
 - 3 or more segmental rib fractures (flail chest)
 - 3 or more bi-cortically displaced/offset rib fractures
 - clinical findings of paradoxical motion
 - instability or "clicking" on palpation of chest wall or as reported by the patient
 - Failure to wean from ventilator

Contraindications

1. Absolute
 - shock/ongoing resuscitation
 - severe traumatic brain injury

- acute myocardial infarction
- fractures outside of ribs 3-10

2. Relative

- Age <18 yrs
- Age >80 yrs
- unstable spine injury
- empyema
- history of chest wall radiation
- mild to moderate traumatic brain injury

Timing


1. Non-ventilated patients

- when feasible, less than 24 hrs is optimal
- should be performed within 72 hours of injury
- SSRF should be delayed in the face of higher priority injuries

2. Ventilated patients

- earliest feasible time for flail indication
- should be performed within 72 hrs of injury for non-flail indications.
- SSRF should be delayed in the face of higher priority injuries.

SSRF ALGORITHM



DEFINITIONS OF TERMS

SEVERE TBI

- ANY GCS <8
- SIGNS OF INTRACRANIAL HYPERTENSION

RELATIVE CONTRAINDICATIONS

- AGE <18 YEARS
- AGE >80 YEARS
- UNSTABLE SPINE INJURY
- EMPYEMA
- PRIOR CHEST WALL RADIATION
- MILD/MODERATE TBI

CHEST WALL INSTABILITY

FLAIL SEGMENT

- 3+ IPSILATERAL CONSECUTIVE RIBS WITH FRACTURES IN 2 LOCATIONS
- CLINICAL FINDING OF PARADOXICAL MOTION

OFFSET FRACTURES

- 3+ IPSILATERAL RIB FRACTURES WITH DISPLACEMENT OF 100% OF RIB WIDTH ON AXIAL CT

INSTABILITY OR "CLICKING" ON PALPATION OR REPORTED BY THE PATIENT

3+ > 50% DISPLACEMENT

- THREE IPSILATERAL CONSECUTIVE OR NON-CONSECUTIVE RIBS EACH WITH A FRACTURE DISPLACED 50% OF THE RIB WIDTH ON AXIAL CT

PULMONARY DERANGEMENTS

- RESPIRATORY RATE >20
- INCENTIVE SPIROMETRY <50% OF PREDICTED
- NUMERICAL PAIN SCORE >5/10
- POOR COUGH

FAILURE TO WEAN

MUST BE CLINICALLY DETERMINED TO BE RELATED TO THE RIB FRACTURES

UNABLE TO PROGRESS TO SPONTANEOUS BREATHING TRIAL AFTER 48 HOURS

ABLE TO OBTAIN SPONTANEOUS BREATHING TRIAL FOR 60 MINUTES BUT DEVELOPS ≥2 OF THE FOLLOWING

- INCREASED RESP. RATE >35
- INCREASED HEART RATE >140
- OXYGEN SATURATION <90%
- RSSI >105
- ANXIETY
- DIAPHORESIS
- AGITATION

OF NOTE: VENTILATOR WEANING SHOULD BE AT THE DISCRETION OF THE TREATING BEDSIDE PHYSICIAN.

HIGHER PRIORITY INJURIES

- PRE-OPERATIVE SPINAL INJURY
- OPEN ABDOMEN
- SIGNIFICANT VASCULAR TRAUMA
- PELVIC EXTERNAL FIXATION

References

1. Bauman, ZM., Grams, B., Yanala, U., Shostrom, V., Waibel, B., Evans, CH., Cemaj, S., & Schlitzkus, L. (2020). Rib fracture displacement worsens over time. *European Journal of Trauma and Emergency Surgery*. 2020 March 27. doi: 10.1007/s00068-020-01353-w
2. Cataneo, AJ., Cataneo, DC., de Oliveria, FH., Arruda, KA., El Dib, R., & de Oliveira Carvalho, PE. (2015). Surgical versus nonsurgical interventions for flail chest. *Cochrane Database of Systematic Reviews*, 2015(7), 1-30. doi: 10.1002/1465158.CD009919.pub2.
3. DeFrees, L., Tafen, M., Bhakta, A., Ata, A., Martone, S., Glotzer, O., Krautsak, K., Rosati, C., Stain, S., & Bonville, D. (2015). Open reduction and internal fixation of rib fractures in polytrauma patients with flail chest. *American Journal of Surgery*, 211(4), 761-767. doi: 10.1016/j.amjsurg.2015.11.014.
4. Kasotakis, G., Hasenboehler, EA., Streib, EW., Patel, N., Patel, MB., Alarcon, L., Bosarge, PL., Love, J., Haut, ER., & Como, JJ. (2017). Operative fixation of rib fractures after blunt trauma: A practice management guideline from the Eastern Association for the Surgery of Trauma. *The Journal of Trauma and Acute Care Surgery*, 82(3), 618-626. doi: 10.1097/TA.0000000000001350
5. Majercik, S., Vijayakumar, S., Olsen, G., Gardner, S., Granger, SR., Van Boerum, DH., & White, TW. (2015). Surgical stabilization of severe rib fractures decreased incidence of retained hemothorax and empyema. *American Journal of Surgery*, 210(8), 1112-1116. doi: 10.1016/j.amjsurg.2015.08.008.
6. Pieracci, FM., Coleman, J., Ali-Osman, F., Mangram, A., Majercik, S., White, TW., Jeremitsky, E., & Doben, AR. (2018). A multicenter evaluation of the optimal timing of surgical stabilization of rib fractures. *Journal of Trauma and Acute Care Surgery*, 84(1), 1-10. doi: 10.1097/TA.0000000000001729
7. Pieracci, FM., Lin, Y., Rodil, M., Synder, M., Herbert, B., Tran, DK., Stoval, RT., Johnson, JL., Biffi, WL., Barnett, CC., Cothren-Burlew, C., Fox, C., Jurkovich, GJ., & Moore, EE. (2018). A prospective, controlled clinical evaluation of surgical stabilization of severe rib fractures. *Journal of Trauma and Acute Care Surgery*, 80(2), 187-194. doi: 10.1097/TA0000000000000925.
8. Pieracci, FM., Majercik, S., Ali-Osman, F., Ang, D., Doben, A., Edwards, JG., French, B., Gasparri, M., Marasco, S., Minshall, C., Sarani, B., Tisol, W., VanBoerum, DH., & White, TW. (2017). Consensus statement: Surgical stabilization of rib fractures rib fracture colloquium clinical practice guidelines. *Internal Journal of the Care of the Injured*, 48(2), 307-321. doi: 10.1016/j.injury.2016.11.026.
9. Witt, C., & Bulger, E. (2017). Comprehensive approach to the management of the patient with multiple rib fractures: a review and introduction of a bundled rib fracture management protocol. *Trauma Surgery & Acute Care Open*, 2(1), 1-7. doi: 10.1136/tsaco-2016-000064.
10. Wada, T., Yasuaga, H., Inokuchi, R., Matsui, H., Matsubara, T., Ueda, Y., Gunshin, M., Ishii, T., Doi, K., Kitsuta, Y., Nakajima, S., Fushimi, K., & Yahagi, N. (2015). Effectiveness of surgical rib fixation on prolonged mechanical ventilation in patients with traumatic rib fractures: A propensity score-matched analysis. *Journal of Critical Care*, 30(6), 1227-231. doi: 10.1016/j.jccr.2015.07.027.

Author(s)

Zachary Bauman, DO

Last Updated

May, 2020

Revision #1

Created 27 June 2023 16:33:00 by Emily Cantrell

Updated 17 October 2024 17:19:32 by Emily Cantrell